

Configurational entropy of hard spheres

Luca Angelani¹ and Giuseppe Foffi²

¹ Research center SMC INFM-CNR, c/o Università di Roma “La Sapienza”, P.le A. Moro 2, I-00185, Roma, Italy

² Institut Romand de Recherche Numérique en Physique des Matériaux IRRMA, and Institute of Theoretical Physics (ITP), Ecole Polytechnique Fédérale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland

Abstract. We numerically calculate the configurational entropy S_{conf} of a binary mixture of hard spheres, by using a perturbed Hamiltonian method trapping the system inside a given state, which requires less assumptions than the previous methods [R.J. Speedy, Mol. Phys. **95**, 169 (1998)]. We find that S_{conf} is a decreasing function of packing fraction φ and extrapolates to zero at the Kauzmann packing fraction $\varphi_K \simeq 0.62$, suggesting the possibility of an ideal glass-transition for hard spheres system. Finally, the Adam-Gibbs relation is found to hold.

1. Introduction

The idea that the glass transition is driven by a decreasing of the number of accessible states upon lowering temperature (or raising density) is quite old [1, 2, 3]. In this picture, if crystallization is avoided, an ideal glass transition is expected to happen at the point where the configurational entropy S_{conf} (the logarithm of the number of states) vanishes. When the liquid enters in the supercooled region, the dynamics becomes slower and slower and the particles get trapped for an increasingly longer time inside the “cages” made by their neighbors: the dynamics of the system can be successfully described as a “fast” motion of the representative point in the $3N$ configuration space inside metastable states, and a “slow” motion corresponding to jumps among states. Entering more in the supercooled region the number of accessible metastable states decreases and the extrapolation to zero of S_{conf} defines the ideal glass transition. In experiments (or numerical simulations) the region close to the ideal glass transition is unreachable, due to the “apparent” arrest of the system at the so called glass-transition temperature (or density) when relaxation times become longer than experimental time scale. The above scenario has been shown to be valid for many interacting systems, based on smooth pair-potential (as Lennard-Jones liquids), for which the Potential Energy Landscape (PEL) approach [4, 5, 6, 7, 8, 9, 10, 11, 12, 13] and the replica method [14, 15] have allowed to give numerical estimations of S_{conf} and of the ideal glass transition.

The overall picture is still not well established for Hard Spheres (HS), for which the existence of a glass transition is still an open question [16, 17, 18, 19, 20]. A particularly

important role seems to be played by the dimensionality of the system. In particular in $d=2$ dimension there are numerical [20, 21] and theoretical [22, 23] evidences of the absence of a thermodynamic glass transition, while the opposite seems to be true in $d=3$ [18, 24]. Moreover, the step-wise form of the interparticle potential does not allow a PEL analysis and different approaches have to be taken in consideration in order to calculate the configurational entropy S_{conf} . In the past, attempts to estimate S_{conf} have been performed based on different evaluations of the entropy in each single state [18, 25, 26]. Recently the replica method has been extended to the HS case for one-component systems [23, 24].

In this paper we follow an approach, based on the Frenkel-Ladd method [27] and recently introduced in the study of Lennard-Jones systems [28] and attractive colloids [29, 30], to numerically estimate S_{conf} for binary hard spheres. As in previous studies, the calculation is reduced to that of vibrational entropy S_{vib} , using the fact that the total entropy S can be decomposed into the sum of a configurational contribution S_{conf} and a vibrational one S_{vib} :

$$S = S_{conf} + S_{vib} . \quad (1)$$

This expression is consistent with the idea that, at high enough density, there are two well separated time scales: a fast one, related to the motion inside a local state (the rattling in the cage), and a slow one associated to the exploration of different states.

The total entropy S is obtained by thermodynamic integration, starting from the ideal gas state. The quantity S_{vib} is calculated using a perturbed Hamiltonian, adding to the original Hamiltonian an harmonic potential around a given reference configuration. Calculating the mean square displacement from the reference configuration and making an integration over the strength of the perturbation, it is possible to estimate the vibrational entropy [29]. The difference $S - S_{vib}$ provides an estimate of the configurational entropy S_{conf} as a function of packing fraction φ (or density ρ).

The main findings of the present work are the following. *i)* S_{conf} is a decreasing function of packing fraction φ , and a suitable extrapolation to zero provides and estimate of the ideal phase transition point (Kauzmann packing fraction) $\varphi_K \simeq 0.62$. *ii)* The diffusivity D and configurational entropy S_{conf} are related through the Adam-Gibbs relation, in agreement with previous claims [18].

2. The model

The studied system is a binary mixture 50–50 of hard spheres, A and B , with diameters ratio $\sigma_B/\sigma_A=1.2$. The collision diameters are $\sigma_{AA}=\sigma_A$, $\sigma_{BB}=\sigma_B$ and $\sigma_{AB}=(\sigma_A + \sigma_B)/2$. The particles ($N=256$) are enclosed in a cubic box with periodic boundary conditions. We use the following units: σ_B for length and $m_A = m_B = 1$ for mass. Moreover we chose $k_B = 1$ and $\hbar = 1$. The density is measured by the packing fraction φ that is related to the number density $\rho = N/V$ by $\varphi=\rho\pi(\sigma_A^3 + \sigma_B^3)/12$. We analyzed systems in the range $\varphi=0.425 - 0.580$. Hard spheres systems depend only trivially on temperature




Figure 1. Mean square displacements (MSD) of the species A (top) and B (bottom). Both MSDs has been normalized by the value of the diameter squared. The dashed lines represent the cage size squared, δr^2 .

that sets an overall scale for the dynamics, consequently we perform all our simulations at $T = 1$. We performed standard event-driven molecular dynamics [31] and we stored several equilibrated configurations at different density.

3. Diffusivity

The diffusion coefficients D of the two species have been extracted from the long time limit of the mean square displacements (MSD) $\langle r^2(t) \rangle = N^{-1} \langle [\mathbf{r}(t) - \mathbf{r}(0)]^2 \rangle$ ($\mathbf{r}$ is the $3N$ -vector of the coordinates):

$$\lim_{t \rightarrow \infty} \frac{\langle r^2(t) \rangle}{t} \simeq 6D. \quad (2)$$

To improve the statistical significance of the data, an average over 10 independent runs have been performed. In Fig 1 the mean squared displacements for the slowest cases, i.e. $\phi > 0.56$, are presented for both species. It is clear that on increasing the density the MSDs develop the typical two step relaxation pattern. The first part of the MSD is purely ballistic while, at later stage, it reaches the diffusive regime, described by Eq. 2. Between this two regimes a plateau starts to develop. This is the clear indication of a caging effect. Each particle starts to feel the crowding of its neighbors and it is trapped in a cage for longer and longer time on increasing the density. The height of the plateau is the typical “cage diameter squared”, δr^2 . For both species we find $\delta r^2 = 3 \cdot 10^{-2} \sigma_\alpha^2$ for $\alpha = A$ or B represented by a dashed line in Fig. 1. This is a clear evidence that the two species have the same caging effect. We shall return on the value of δr^2 later on.

In Fig. 2 the diffusivities D of A and B particles are plotted as a function of the packing fraction ϕ . Dashed lines in the figure are power-law fits $D = C (\phi_c - \phi)^\gamma$




Figure 2. Diffusivity for A and B particles as a function of packing fraction ϕ . The lines are power-law fits for $\phi \geq 0.53$, $C (\phi_c - \phi)^\gamma$, with: $\phi_c=0.583$, $\gamma=2.27$, $C=9.50$ for A particles; $\phi_c=0.583$, $\gamma=2.47$, $C=11.66$ for B particles.

of the high packing-fraction data ($\phi \geq 0.53$), as predicted by mode-coupling theory. The fitted parameters are: $\phi_c=0.583$, $\gamma=2.27$, $C=9.50$ for A particles and $\phi_c=0.583$, $\gamma=2.47$, $C=11.66$ for B particles. We note that both diffusivities give rise to the same mode-coupling packing fraction ϕ_c , in agreement with the prediction of the theory [32] and with previous simulations of the same model [33].

4. Configurational entropy

We now turn to the calculation of configurational entropy. The method we follow to estimate S_{conf} requires the computation of the total entropy S and vibrational entropy S_{vib} . The total entropy S is calculated *via* a thermodynamic integration from ideal gas and can be expressed as

$$S(\rho) = S_{id}(\rho) + S_{ex}(\rho) , \quad (3)$$

where S_{id} is the entropy of the ideal gas and S_{ex} is the excess entropy with respect to the ideal gas. For a binary mixture, the ideal gas entropy is:

$$\frac{S_{id}(\rho)}{N} = \frac{5}{2} - \ln \rho - 3 \ln \lambda + \ln 2 , \quad (4)$$

where $\lambda = (2\pi\beta\hbar^2/m)^{\frac{1}{2}}$ is the De Broglie wavelength ($\hbar$ is the Planck's constant and has been set to unitary value), and the term $\ln 2$ takes into account the mixing contribution. The term S_{ex} can be expressed in the following form

$$S_{ex}(\rho) = -\frac{N}{T} \int_0^\rho \frac{d\rho}{\rho^2} P_{ex} , \quad (5)$$




Figure 3. Excess entropy S_{ex} for the mixture of hard spheres as obtained from our simulations (symbols) compared with the analytical Carnahan-Starling expression (line) [35, 36].

with P_{ex} the excess pressure. We extracted P_{ex} from the zero density limit up to the densities of interest, performing numerical simulations and fitting the results of the pressure with a high order polynomial in ρ . In Fig. 3 we show the numerically calculated excess entropy S_{ex} (symbols) together with the analytic estimate provided by the Carnahan-Starling (CS) equation of state, extended to hard sphere mixtures [34, 35, 36]. We note that at high densities the CS equation of state overestimates the entropy of about 7%. This discrepancy, however, is not sufficiently significant to affect the resulting S_{conf} values, in particularly closed to the glass transition.

The method we use for the calculation of S_{vib} is based on the investigation of a perturbed system

$$\beta H' = \beta H + \alpha N(\mathbf{r} - \mathbf{r}_0)^2, \quad (6)$$

where H is the unperturbed hard spheres Hamiltonian, α is the strength of the perturbation, $\mathbf{r}_0$ specifies the particles coordinates of a reference configuration and $(\mathbf{r} - \mathbf{r}_0)^2 \equiv N^{-1} \sum_{i=1}^N (\vec{r}_i - \vec{r}_{0,i})^2$. The reference configuration $\mathbf{r}_0$ is chosen from equilibrium configurations at the considered density (randomly extracted from the stored configurations obtained during molecular dynamics simulations). With this choice one is sure that the estimated vibrational entropy (see formula below) pertains to the correct state at the studied density. The vibrational entropy can be obtained from the formula (see Ref. [29] for details):

$$\frac{S_{vib}}{N} = \int_{\alpha_0}^{\alpha_\infty} d\alpha' \langle (\mathbf{r} - \mathbf{r}_0)^2 \rangle_{\alpha'} - \frac{3}{2} \ln \left(\frac{\alpha_\infty \lambda^2}{\pi} \right) + \frac{3}{2}, \quad (7)$$

where $\alpha_{0,\infty}$ are the lower/upper limit of integration, and $\langle \dots \rangle_{\alpha'}$ is the canonical average for a given α' . The choice of α_0 deserves few comments. If the system were confined




Figure 4. The quantity $\langle(\mathbf{r} - \mathbf{r}_0)^2\rangle_\alpha$ plotted vs. α in logarithmic scale for different packing fractions φ . Vertical lines are the values $\alpha_0^{(1)} = 2^{2.5}$ and $\alpha_0^{(2)} = 2^{4.5}$ used as α_0 -value in the integral in Eq. 7 for the calculation of S_{vib} .

to move inside a given local free-energy minimum, for a correct estimation of S_{vib} one would take the lower limit $\alpha_0=0$ in the integral in Eq. 7. As the system, at enough low value of α , begins to sample different states (the harmonic force due to the perturbation is no more able to constrain the system inside one state), α_0 has to be chosen in such a way that the system has not yet left the state: the underlying idea is that Eq. 7 gives a correct estimation of S_{vib} until the system remains trapped in the state. An appropriate choice in our case seems to be $\alpha_0^{(1)}=2^{2.5}$ for all the densities, as, close to this point, one observes a crossover for all the investigated densities (more pronounced for low density data). In Fig. 4 the quantity $\langle(\mathbf{r} - \mathbf{r}_0)^2\rangle_\alpha$ is reported as a function of α . An arrow indicates the chosen value $\alpha_0=\alpha_0^{(1)}$, below which one observes the crossover associated to the exploration of different states.

It is worth noting that different choices of α_0 are in principle possible, giving rise to different estimations of the vibrational entropy term. However, even though a kind of arbitrariness is present in the method, one can argue that a reasonable choice should be for values above $\alpha_0^{(1)}$ (close to the crossover corresponding to the exploration of different states) and below an upper value $\alpha_0^{(2)}$ at which one is sure that the system is still confined in a single state. The latter value can be estimated requiring that the MSD $\langle(\mathbf{r} - \mathbf{r}_0)^2\rangle$ is always close/below the cage diameter squared $\delta r^2 \simeq 3 \cdot 10^{-2} \sigma_\alpha^2$ (with $\alpha = A$ or B) (this has been estimated from the *plateau* of the mean square displacement, see Fig. 1).

The chosen value in our case is $\alpha_0^{(2)}=2^{4.5}$ (indicated by an arrow in Fig. 4). We then repeated the same calculation of S_{vib} using Eq.7 with the lower bound in the integral $\alpha_0 = \alpha_0^{(2)}$. In this way we obtain a lower and upper bound for the quantity of interest S_{vib} , by using respectively $\alpha_0^{(1)}$ or $\alpha_0^{(2)}$ in the expression of S_{vib} in Eq.7.




Figure 5. Configurational entropy S_{conf} as function of packing fraction φ . Open symbols are data using $\alpha_0^{(1)} = 2^{2.5}$, full symbols using $\alpha_0^{(2)} = 2^{4.5}$ (see text). Dashed and dot-dashed lines are from Speedy [18] for binary and monatomic hard-spheres respectively. Thin full line is the analytical computation of Parisi and Zamponi for monatomic hard-spheres [24]. Thick lines are polynomial extrapolations of our data in the high packing fraction region, giving rise to the same Kauzmann packing fraction estimation for which $S_{conf}(\varphi_K)=0$: $\varphi_K \simeq 0.62$.

Fig. 5 shows S_{conf} as a function of φ . The configurational entropy S_{conf} is calculated using Eq. 1, where the two entropies S and S_{vib} are obtained from Eq.s 3 and 7 respectively. Due to the fact that the correct integral for the estimation of S_{vib} should be done from $\alpha_0=0$, but with the system always inside a given state, we have added to the expression in Eq. 7 the term $\alpha_0 \langle (\mathbf{r} - \mathbf{r}_0)^2 \rangle_{\alpha_0}$, corresponding to assume a constant value of $\langle (\mathbf{r} - \mathbf{r}_0)^2 \rangle$ below α_0 and using a zero value for the lower limit of the integral in Eq. 7. Fig. 5 shows the two estimates of S_{conf} , corresponding to the two different values of α_0 : $\alpha_0^{(1)}=2^{2.5}$ (open symbols) and $\alpha_0^{(2)}=2^{4.5}$ (full symbols). One observes that the discrepancy between the two estimations decreases by increasing the packing fraction, suggesting that, at high density, the method used to calculate S_{conf} is less affected by the choice of the free parameters entering in its evaluation. This is probably due to the fact that increasing the density the system tends to be more trapped in a local free-energy minimum. Indeed, it is only at high density that the method is expected to work better, due to the better definition of two time scales corresponding to local-fast and global-slow dynamics (see Fig. 1). At low density, instead, the two are less separated and this corresponds to a difficulty in the extrapolation for $\alpha_0 \rightarrow 0$ of the quantity reported in Fig. 4. The low density data show a more clear crossover on lowering α , and then a worst definition of state in this limit. As we are interested in the high packing fraction extrapolation, this fact do not affect our prediction on the Kauzmann density value. Also reported in the figure are the curves obtained by




Figure 6. The Adam-Gibbs relation $D=D_{\infty} \exp[-N\Delta/(TS_{conf})]$ ($T=1$) for the two species A and B: $D_{\infty}=3.66$, $\Delta=9.8$ for A particles; $D_{\infty}=4.59$, $\Delta=10.5$ for B particles. The data of S_{conf} are obtained with $\alpha_0=\alpha_0^{(2)}$.

Speedy [18] using a different method (assuming a particular form of the vibrational entropy, a Gaussian distribution of states and involving some free parameters) for the estimation of S_{conf} , for monatomic (dot-dashed line) and binary (dashed line) hard-spheres (with the same diameters ratio 1.2 and composition 50 : 50 as in our case). It is worth noting that our method improves on Speedy's one, as, even though requiring some accuracy in the choice of the α_0 parameter, has the advantage to be less affected by the presence of many free parameters and particular assumptions. We note that the data of Speedy for the binary case do agree very well with our data with $\alpha_0=\alpha_0^{(2)}$, suggesting the possibility that the choice of $\alpha_0=\alpha_0^{(2)}$ is more accurate for the estimation of S_{vib} and so of S_{conf} . As a comparison, in Fig. 5 is also reported an analytic estimation of S_{conf} recently provided by Parisi and Zamponi [24] for monatomic hard-spheres. From the φ -dependence of the configurational entropy one can determine the packing fraction at which S_{conf} extrapolates to zero, corresponding to the ideal phase transition point (Kauzmann packing fraction φ_K) $S_{conf}(\varphi_K)=0$. Using a polynomial extrapolation [37] for the two set of data (corresponding to the different α_0 values) we obtain an estimated Kauzmann packing fraction value $\varphi_K \simeq 0.62$ (see Fig.5). It is worth noting that, even though the two curves are quite different, the estimated value of φ_K is the same, again suggesting the robustness of the method in the high density region and then in the estimation of the Kauzmann packing fraction.

5. Adam-Gibbs relation

In this Section we explore the validity of the Adam-Gibbs relation, linking dynamic quantities, like diffusivity, to S_{conf} . In Fig. 6 we report the diffusivities D for A and B particles vs. the quantity $(TS_{conf})^{-1}$, with S_{conf} obtained for the value $\alpha_0 = \alpha_0^{(2)}$. We find that the Adam-Gibbs relation

$$D = D_\infty e^{-N\Delta/TS_{conf}} \quad (8)$$

is verified (lines in the figure), with: $D_\infty = 3.66$, $\Delta = 9.8$ for A particles; $D_\infty = 4.59$, $\Delta = 10.8$ for B particles. A similar behavior is obtained using S_{conf} calculated with $\alpha_0 = \alpha_0^{(1)}$ (not shown in the figure), with the values: $D_\infty = 24.5$, $\Delta = 9.5$ for A particles; $D_\infty = 37.5$, $\Delta = 10.5$ for B particles, suggesting that, in this range of diffusivity values, the AG expression is not able to discriminate between the two different estimations of S_{conf} .

6. Conclusions

In conclusion, we have calculated S_{conf} for binary mixture hard spheres, by numerically estimating the total entropy S (*via* thermodynamic integration from ideal gas) and the vibrational entropy S_{vib} using a numerical procedure based on Frenkel-Ladd method and recently applied in the analysis of Lennard-Jones systems and attractive colloids: the system is constrained inside a given “*state*” through an harmonic perturbed term in the Hamiltonian. We found, in agreement with analytical and simulation results in the literature, that S_{conf} is a decreasing function of the packing fraction φ , suggesting the possibility of a vanishing of S_{conf} around the Kauzmann point $\varphi_K = 0.62$. Moreover, by studying the relationship between S_{conf} and the diffusion constant D , the Adam-Gibbs relation is found to reasonably hold for the analyzed system.

Acknowledgments

We thank G Ruocco, F Sciortino and F Zamponi for useful discussions and suggestions. G F acknowledges the support of the Swiss Science Foundation (Grant No. 200021-105382/1).

References

- [1] Kauzmann A W 1948 *Chem. Rev.* **43** 219
- [2] Adam G and Gibbs J H 1958 *J. Chem. Phys.* **43** 139
- [3] Gibbs J H and Di Marzio E A 1958 *J. Chem. Phys.* **28** 373
- [4] Debenedetti P G and Stillinger F H 2001 *Nature* **410** 259
- [5] Stillinger F H and Weber T A 1984 *Science* **225** 983
- [6] Stillinger F H 1995 *Science* **267** 1935
- [7] Sciortino F, Kob W and Tartaglia P 1999 *Phys. Rev. Lett.* **83** 3214
- [8] Scala A, Starr F W, La Nave E, Sciortino F and Stanley H E 2000 *Nature* **406** 166
- [9] Sastry S, Debenedetti P G and Stillinger F H 1998 *Nature* **393** 554
- [10] Büchner S and Heuer A 2000 *Phys. Rev. Lett.* **84** 2168

- [11] Sastry S 2001 *Nature* **409** 164
- [12] Keyes T 1997 *J. Phys. Chem.* **101** 2921
- [13] Sciortino F 2005 *J. Stat. Mech.* P05015
- [14] Mézard M and Parisi G 1999 *J. Chem. Phys.* **111** 1076
- [15] Coluzzi B , Parisi G and Verrocchio P 2000 *J. Chem. Phys.* **112** 2933
- [16] Rintoul M D and Torquato S 1996 *Phys. Rev. Lett.* **77** 4198
Rintoul M D and Torquato S 1996 *J. Chem. Phys.* **105** 9258
- [17] Robles M, Lòpez de Haro M, Santos A and Bravo Yuste S 1998 *J. Chem. Phys.* **108** 1290
- [18] Speedy R J 1998 *Mol. Phys.* **95** 169
- [19] Tarzia M, de Candia A, Fierro A, Nicodemi M and Coniglio A 2004 *Europhys. Lett.* **66** 531
- [20] Donev A, Stillinger F H and Torquato S 2006 *Phys. Rev. Lett.* **96** 225502
- [21] Santen L and Krauth W 2000 *Nature* **405** 550
- [22] Tarzia M 2007 *J. Stat. Mech.* P01010
- [23] Zamponi F 2006 *Phil. Mag.* **87** 485
- [24] Parisi G and Zamponi F 2005 *J. Chem. Phys.* **123** 133501
- [25] Speedy R J 1999 *J. Chem. Phys.* **110** 4559
Speedy R J 2001 *J. Chem. Phys.* **114** 9069
- [26] Aste T and Coniglio A 2003 *Physica A* **330** 189
Aste T and Coniglio A 2003 *J. Phys.: Condens. Matter* **15** S803
Aste T and Coniglio A 2004 *Europhys. Lett.* **67** 165
- [27] Frenkel D and Smit B 2001 *Understanding Molecular Simulation* (Academic Press, London, 2nd)
- [28] Coluzzi B, Mézard M, Parisi G, and Verrocchio P 1999 *J. Chem. Phys.* **111** 9039
- [29] Angelani L, Foffi G, Sciortino F and Tartaglia P 2005 *J. Phys.: Condens. Matter* **17** L113; Foffi G and Angelani L (in preparation)
- [30] Moreno A J, Buldyrev S V, La Nave E, Saika-Voivod I, Sciortino F, Tartaglia P and Zaccarelli E 2005 *Phys. Rev. Lett.* **95** 157802
- [31] Rapaport D C 1997 *The art of computer simulations* (Cambridge Univ Press, London, 2nd)
- [32] Voigtmann Th 2003 *Phys. Rev. E* **68** 051401
- [33] Foffi G, Götze W, Sciortino F, Tartaglia P and Voigtmann Th 2003 *Phys. Rev. Lett.* **91** 085701
Foffi G, Götze W, Sciortino F, Tartaglia P and Voigtmann Th 2004 *Phys. Rev. E* **69** 011505
- [34] Alder B J 1964 *J. Chem. Phys.* **40** 2724
- [35] Mansoori G A, Carnahan N F, Starling K E, and Leland T W Jr 1971 *J. Chem. Phys.* **54** 1523
- [36] The analytical CS expression for S_{ex} is given by Eq. 9 in Ref. [35] omitting the last term $\ln Z$, as in our case the excess entropy is defined with respect to the ideal gas at the same density instead of at the same pressure as in Ref. [35]
- [37] We use a polynomial function of the form $A(\varphi_K - \varphi) - B(\varphi_K - \varphi)^2$ to fit the data with $\varphi > 0.5$. Adding more polynomial orders do not significantly affect the extrapolated Kauzmann packing fraction value.